

Article

Influence of Alcohol Consumption, Personality and Attention Deficit Hyperactivity Disorder on Traffic Offenders

Cristina Escamilla-Robla¹, M. Ángeles Beleña-Mateo² and Joaquín Mateu-Mollá³

1 Universidad Europea de Valencia.

2 Universidad de Valencia.

3 Universidad Internacional de Valencia.

ARTICLE INFO

Received: January 22, 2022

Accepted: April 20, 2022

Keywords:

Road crime
Prevention
Personality
Alcohol abuse
Attention deficit hyperactivity disorder

ABSTRACT

Background: Despite the authorities' efforts to promote responsible driving, traffic accidents are a notable cause of death in many countries. Scientific research shows that variables such as alcohol abuse and personality dimensions can contribute to explaining this social phenomenon and the rising number of traffic offences. The present study aims to explore how alcohol abuse, personality disorders and Attention Deficit Hyperactivity Disorder (ADHD) are associated with risky driving behaviour. **Method:** the sample included 300 subjects: the study group (199 convicted of traffic offences) and the control group (101 subjects from the general driving population). The tools used were the International Personality Disorder Scale, the Alcohol Use Disorders Identification Test and the Adult ADHD Self-Report Scale (ASRS). **Results:** a higher incidence of signs of paranoid, antisocial, impulsive, borderline and anxious personality traits was detected for the study group, as well as a higher prevalence of alcohol abuse. **Conclusions:** this research offers indications for the design of effective preventive and reinsertion interventions. It also highlights the importance of considering psychological dimensions to ensure road safety.

Influencia del Consumo de Alcohol, la Personalidad y el Trastorno de Déficit de Atención e Hiperactividad en Delincuentes viales

RESUMEN

Antecedentes: Los accidentes de tráfico son una causa de muerte en muchos países, pese a los esfuerzos por promover una conducción responsable. La investigación sobre el problema muestra que variables como el abuso de alcohol y dimensiones de personalidad, contribuyen a explicar este fenómeno social y el aumento de la delincuencia vial. El propósito de este estudio es explorar cómo el abuso de alcohol, los trastornos de la personalidad y el Trastorno por Déficit de Atención e Hiperactividad (TDHA) se asocian a conductas de riesgo al volante. **Método:** Participaron 300 sujetos: un grupo de estudio (199 personas condenadas por delitos viales) y otro control (101 sujetos extraídos de la población general de conductores). Los instrumentos utilizados fueron el examen internacional de trastornos de la personalidad (IPDE), el cuestionario de identificación de trastornos por consumo de alcohol (AUDIT) y el cuestionario autoinformado de cribado del TDAH. **Resultados:** el grupo estudio presentó mayor incidencia de indicios de trastornos de personalidad paranoide, disocial, impulsivo, límite y ansioso; así como mayor prevalencia de abuso de alcohol. **Conclusiones:** El estudio ofrece claves para construir intervenciones preventivas y de reinsertión eficaces, enfatizando la relevancia de dimensiones psicológicas para garantizar la seguridad en las vías públicas.

Palabras clave:

Delincuencia vial
Prevención
Personalidad
Abuso de alcohol
Trastorno por déficit de atención e hiperactividad

Road traffic accidents are one of the main causes of death for the population aged 5-29. Not only do they pose a serious public health problem despite progress being made (World Health Organization [WHO], 2018), but they also entail very high social costs (Wijnen & Stipdonk, 2016).

Around 22,800 people died in road traffic accidents in EU Member States in 2019 (European Commission, 2020). About 25% of these deaths could have been avoided, since alcohol was the main reason for these accidents in 2018 (Le Lièvre et al., 2019) and in 2016 (Calinescu & Adminaite, 2018).

Spain registered 104,080 accidents involving people in 2019, in which 1,755 died and 8,613 required hospitalization. In terms of risk behaviours, alcohol consumption appeared as the second most common concurrent factor (in 24% of the aforementioned fatal accidents) after distractions (Directorate-General for Traffic, 2020). Moreover, the total number of accidents involving people increased slightly in 2019, rising by 1,781 compared to 2018. This indicates that the behaviour of drivers is still risky. According to Spanish citizens, these accidents are also one of the most important social problems, after cancer and terrorism (Alonso et al., 2019).

The Spanish National Institute of Toxicology and Forensic Sciences (2020) indicates that the toxicological analyses performed with *post-mortem* samples to determine the alcohol, drugs of abuse and psychodrugs used by drivers who died in 2019 revealed that 45.50% of them obtained positive results. This tendency has grown in recent years. In 61.80% of these cases, alcohol was detected either on its own or in combination with other substances and, of these, 81.50% had blood alcohol levels of > 1.2 g/L (the threshold for prosecution).

Apart from the increase in road accident deaths in recent years, convictions for road safety crimes totalled 80,480 in 2019 (General State Prosecutor's Office, 2020) in addition to 24,001 community work sentences. Of these sentences, 60% involved driving while under the influence of alcohol or other drugs, which are the offences described in Article 379.2 (Spanish Organic Law 15/2007) and the most relevant typology of those considered (Alonso et al., 2015). It is worth reflecting on these in relation to the political-criminal measures taken in recent years.

In the EU directive On Driving Licences (Directive 2006/126/EC) one of the proposals for Member States was to set a number of minimum requirements for issuing driving licenses. For instance, drivers must "take account of all the factors affecting driving behaviour (alcohol, fatigue, poor eyesight, etc.) so as to retain full use of the faculties needed to drive safely" (p. 46). In Spain, the Spanish Royal Decree of 8 May (RD 818/2009), on the General Drivers Regulation, put into practice Directive 2006/126/EC which establishes the psycho-physical capability tests to verify that no disease or disorder is present that may render drivers incapable of driving, such as personality disorders (PDs), Attention Deficit Hyperactivity Disorder (ADHD), or disorders related to substance addiction. Article 36 therefore stipulates that drivers may have their driving licence revoked if the psycho-physical capacity required for safe driving has disappeared or worsened.

Since road accidents or road crimes are not concerned with the awareness or lack thereof of regulations, but rather the behaviour of drivers, personality may be an influential variable (Martí-Belda et al., 2019). Personality is understood as a "persisting pattern

of the way one's environment is perceived, related and devised, and is manifested in different personal and social contexts" (De Miguel & Pelechano, 2000, p. 26). Personality is, therefore, considered an important variable for analysing and understanding the stability, perseverance, and recidivism of criminals' antisocial behaviour, due to the wide range of interindividual variety that exists in such conduct (Gallardo & Andrés, 2009). Moreover, political and sociodemographic conditions may contribute to rates of recidivism (Chen & Jou, 2018).

In terms of road traffic, driving is yet another daily task that takes place in a shared social context (Nordfjaen & Simsekoğlu, 2014). Within this context, personality traits are extremely relevant when studying drivers' behaviour and its relation to accidents or dangerous conduct (Constantinou et al., 2011; Haerani et al., 2014; Jonah et al., 2001; Martí-Belda et al., 2019; Tortosa & Montoro, 2002).

In behavioural terms, it is also necessary to study the existence of PDs, which occur when personality traits are inflexible/non-adaptive in the subject's culture and remain stable over time, harming or causing discontent to the person and his/her environment (Alavi et al., 2017). The American Psychiatric Association's DSM-5 (APA, 2013) indicates that the "general pattern of scorn and violating others' rights" (p. 659) is a determining characteristic of antisocial PDs. Some characteristics of different PDs are associated with other actions that can be considered a threat against safe driving, particularly those expressing antisocial conduct or those that pose a safety risk for people (Alonso et al., 2007; Räisänen et al., 2019; Ramos-Quiroga et al., 2013). This is also set out by Spanish Royal Decree RD 818/2009, where moderate or severe cases of dissocial personality disorder or other disturbers are not admitted, as they are normally accompanied by aggressive behaviour or serious violations of the regulations, negatively affecting road safety.

Cavaola et al. (2003) found higher scores for traffic offenders in behaviours related to psychotic deviation for antisocial behaviours (Gallardo & Andrés, 2009), and higher levels for physical aggression and antisocial practices (Shechory et al., 2011). Hilterman et al. (2012) observed a closer relationship between general psychological problems and those related to alcohol in a group of people convicted of road crimes and serving time in prison, compared to subjects convicted of other crimes. Another study performed with individuals convicted of traffic crimes highlighted high levels of impulsiveness (Bıçaksız & Özkan, 2016), aggressiveness, and neuroticism (Slavinskienė et al., 2015).

Another disorder related to risky or antisocial driving is ADHD, often underestimated and untreated in adults (Eensoo et al., 2015). It is defined by having attention deficit problems, lacking organisation, and hyperactivity-impulsiveness that may persist until adulthood, which consequently results in deteriorated social, academic, and occupational skills (Simon et al., 2009). Different studies relate ADHD to antisocial and other behavioural disorders (Kaye et al., 2014; Klein et al., 2012; López et al., 2005), to an increase in the probability of committing criminal acts (Pérez et al., 2015; Retz et al., 2021), and to higher probability of divorce, unemployment, legal problems, or traffic accidents (Chang et al., 2017; Curry et al., 2017; García et al., 2019; Harpin, 2005; Valero et al., 2017). Drivers with ADHD are more likely to perform dangerous manoeuvres behind the wheel (Dekkers et

al. (2016), to drive without a licence or under the influence of substances, and to commit other traffic violations. These drivers are more prone to suffer accidents compared to the general population (Vaa, 2013). All these behaviours are motivated by impulsiveness and attention deficit (Andreu et al., 2015). ADHD is also associated with disorders caused by taking substances (Kaye et al., 2014; Luderer et al., 2021), making it difficult to concentrate or causing mistakes, which can have serious social and personal impacts (Cabasés & Quintero, 2005; Jiménez-Arriero et al., 2005; Ramos-Quiroga et al., 2006; Valero et al., 2017).

Alcohol consumption seriously alters drivers' capacity to drive safely and increases the risk of having an accident (Alonso et al., 2015; Alonso et al., 2017; Begg et al., 2017; Del Río et al., 2002; Moskowitz & Fiorentino, 2000). It is also more common in drivers whose licence has been revoked (Valero et al., 2017). Almost 20% of drivers attending medical examination centres to re-apply for a driving licence suffer disorders caused by substance use (Gómez-Talegón et al., 2008). Many drivers previously convicted of drink driving tend to be persistent offenders (Monrás et al., 2011; Schell et al., 2006), and alcohol consumption is the best predictor (Gugliotta, 2018; Nelson et al., 2019). Herraiz (2009) and Herraiz et al. (2011) highlighted that 50% of jailed traffic offenders have a drinking problem (Hilterman et al., 2012) and exhibit ethyl alcohol dependence symptoms (Faílde-Garrido et al., 2016; McCutcheon et al., 2009; Valero et al., 2017). Traffic offenders (vs. a control group) drink statistically more alcohol (González-Iglesias & Gómez-Fraguela, 2010). Quite frequently convicted individuals suffer from alcoholism, but only a few are diagnosed, and even those who drive under the influence of alcohol are more likely to relapse and display antisocial behaviour (Freeman et al., 2011; Jornet-Gibert et al., 2013; Keating et al., 2019; Nelson et al., 2015).

Del Río et al. (2001) worked on a study with 8,043 drivers who went to health centres for medical and psychological check-ups to re-apply for their driving licences. These authors observed how 60.30% of them drink alcohol on a regular basis, 7.30% present a problematic alcohol use pattern, and 2% meet alcohol abuse/dependence criteria. Their study revealed that of all those with alcohol consumption problems, 23.20% had been involved in traffic accidents and 18.70% had a record of such offences. Nonetheless, the tests performed with 72.20% of these drivers revealed that they were eligible to drive. These conclusions are not in agreement with the regulations for being granted a driving licence.

The research carried out by Gómez-Talegón et al. (2008) included a sample of 5,234 drivers who went to medical-psychological health centres, of whom 3.30% had mental or substance-related disorders, 18.70% used substances, and 11.70% had impulse-control disorder. Of all these drivers, 39.80% were assessed as "eligible" (most with impulse-control disorder), 53.20% as "eligible, but with restrictions" and 7% as "not eligible". In this last group, 25% suffered a drug abuse or dependence disorder, and 14.30% had dementia. The research carried out by De las Cuevas and Sanz (2009) focused on the effect of mental disorders and psycho-pharmacological treatments on cognitive function and psychomotor skills in driving. It included a sample of patients with mental disease, treated as outpatients, who have a driving licence and drive regularly. Only 20% of them passed

the driving licence tests, and only 10% informed health centres that their driving capacity may be worsening to a certain extent. Of all the subjects, 56.70% did not pass the decision-making test for a mean reaction time, and 22% had a tendency to break the rules. Trujillo et al. (2007) evidenced increased severity of antisocial conduct when antisocial disorder and ADHD occur together, resulting in a more than 50% probability of suffering a behavioural disorder related to rule-breaking (Andreu et al., 2015; Soutullo & Díez, 2007). McDonald and Davey (1996) reviewed studies about the relationship between psychiatric disorders and traffic accidents. They reported that the combination of an antisocial PD and alcoholism increases the vulnerability to accidents. A study performed with drivers with no road crime records, and others convicted for driving under the influence of alcohol, observed that the latter were considerably more likely to adopt antisocial behaviours (Jornet-Gibert et al., 2013).

Lapham et al. (2001) concluded that the road offenders who drive under the influence of alcohol need to be evaluated and treated not only for their potential problems related to risky abuse, but also for the psychiatric disorders that frequently come with alcohol abuse, including, among others, an antisocial personality.

All these data suggest the need to conduct research with a view to modifying current medical examination protocols and tests to increase road safety and to minimise driving under non-optimal conditions, particularly under the effect of psychotropic substances that could alter attention and other processes, as suggested by scholars such as Ayinde et al. (2019). Many arguments prove that it is necessary to build crime policies in relation to drinking, driving and antisocial personality, in order to improve already-existing programmes and to create new evidence-based therapies.

By taking psycho-physical capabilities for driving safely into account, the present study therefore aims to explore those variables related to normative demands in sentenced road traffic offenders (such as harmful alcohol consumption, PDs and attention and impulse-related disorders), and to compare their scores with a control group from the general driving population with no road crime record.

It could be hypothesised that a significant number of road offenders will present signs of harmful alcohol consumption (Faílde-Garrido et al., 2016; González-Iglesias and Gómez-Fraguela, 2010), personality disorders (Kieling et al., 2011; Shechory et al., 2011), and ADHD (Ramos-Quiroga et al., 2006; Valdizán and Izaguerri-Gracia, 2009).

Method

Participants

The sample was made up of 300 participants divided into two groups: a study group with 199 road traffic offenders sentenced to perform community work related to the type of crime (the TASEVAL resocialisation workshop), and a control group made up of 101 subjects with no road crime record who were randomly selected from the general driving population. The inclusion criteria for both groups were being at least 18 years old, while the study group had to attend the aforementioned workshop for the first time, and the control group members had to have a driving licence, with full driving points (according to the Spanish driving point systems),

and have no road crime record. The participants' age ranged from 19 to 76 years of age ($M = 38.47$; $SD = 11.53$ years). Of the whole sample, 78% were male ($n = 234$) and 22% were female ($n = 66$). The sociodemographic characteristics of the sample, divided by groups, can be found in Table 1.

Table 1.
Sociodemographic data of the participants by groups ($N = 300$).

Sociodemographic	Traffic offenders (n = 199)	Non-offender drivers (n = 101)	Total sample (N = 300)
Age (years): M (SD)	36.84 (11.33)	41.69 (11.29)	38.47 (11.53)
Gender: n (%)			
Men	174 (87.40)	60 (59.40)	234 (78.00)
Women	25 (12.60)	41 (40.60)	66 (22.00)
Civil status n (%)			
Single	127 (63.80)	38 (37.60)	165 (55.00)
With couple	72 (36.20)	63 (62.40)	135 (45.00)
Children (yes) n (%)	99 (49.70)	60 (59.40)	159 (53.00)
Education level n (%)			
Primary education / without studies	35 (17.60)	7 (6.90)	42 (14.00)
Secondary education	64 (32.20)	8 (7.90)	72 (24.00)
High school/T&D	57 (28.60)	41 (40.60)	98 (32.70)
University Studies	43 (21.60)	45 (44.60)	88 (29.30)
Employment n (%)			
Student	12 (6.00)	6 (5.90)	18 (6.00)
Unemployed	67 (33.70)	12 (11.90)	79 (26.30)
Active worker	108 (54.30)	78 (77.20)	186 (62.00)
Retired or pensioner	12 (6.00)	5 (5.00)	17 (5.70)

Instruments

International Personality Disorder Examination (IPDE) (Loranger et al., 1997) was used to track subjects who are quite likely to present PDs according to the International Disease Classification (IDC-10; WHO, 1992). The version used (De Miguel & Pelechano, 2000) explores nine PDs (paranoid, schizoid, antisocial, impulsive, borderline, histrionic, anankastic, anxious, dependent) and contains 59 items with dichotomic responses (true/false) that describe behaviour in the past five years. If the subject meets three scale criteria or more (categorical or clinical approach), it suggests that some PDs might be present and the need to later hold an interview to ensure proper diagnosis, if necessary (De Miguel & Pelechano, 2000). This questionnaire's concordance is almost perfect ($Kappa = .87$) (Landis & Koch, 1977).

Alcohol Use Disorders Identification Test (AUDIT) (WHO, 2001). This consists of 10 Likert-type items, with an ordinal response option that assesses alcohol consumption and conduct problems arising from alcohol abuse or dependence (Rubio et al., 1998). Its purpose is to evaluate an at-risk drinker by distinguishing three different levels: alcohol intake, potential dependence on alcohol, and experience of alcohol-related harm. For problematic alcohol consumption, different cut-off points were used for each gender, namely a minimum of 6 points for females and 8 points for males (Péruña de Torres et al., 2005). Therefore, based on these scores, it could be considered that a subject is a high-risk drinker or engages in problematic alcohol use

(Péruña de Torres et al., 2005). In this study, the internal consistency, measured by Cronbach's alpha, was $\alpha = .84$. Other studies obtained similar scores ($\alpha = .82$) (Ballester et al., 2021).

Adult ADHD Self-Report Scale Version 1.1 (WHO, 2003). The screening version by the WHO and Kessler et al. (2005) comprises six items with five Likert-type response options that aim to detect suggestive signs of ADHD. The reference criterion is based on obtaining a score of more than 12 points (Kessler et al., 2007). The first four items assess attention deficit symptoms and the last two explore hyperactivity symptoms. A categorial approach was followed in order to identify subjects with signs of ADHD by calculating the sum of the total score and taking a cut-off point of 12. Specific attention and hyperactivity factors were calculated, too. The reliability of the complete test was adequate in the present study ($\alpha = .68$), measured by Cronbach's Alpha. When the internal consistency for its two factors were independently calculated, it was observed that attention obtained an $\alpha = .74$ and hyperactivity an $\alpha = .60$. These results are very similar to those obtained previously by other authors (Kessler et al., 2007), who found a general reliability of $\alpha = .72$. According to its bifactorial solution, an $\alpha = .82$ was found for attention and an $\alpha = .52$ for hyperactivity (Daigre et al., 2009).

Procedure

After applying for the corresponding authorisation from the Subdepartment of Institutional Relations and Territorial Coordination of the General Secretariat for Penitentiary Institutions to conduct this research work and to obtain its approval, the TASEVAL workshop was chosen (beginning within a period of three months in Valencia, Alicante and Castellón as part of the Valencian Community's Services for Managing Offences and Alternative Penal Measures). Questionnaires were handed out while the workshop was held. For the control group, the subjects who went to health centres to request or re-apply for a driving licence were selected. Before data collection, it was ensured that the participants met the criteria for ethical information, confidentiality, and data processing rules. Informed consents were obtained, duly completed and signed by the participants.

Data analysis

The independent variables were gender, age, ADHD, problems with alcohol consumption, and PDs.

Descriptive analyses were performed using frequencies for the categorical variables, and the frequencies homogeneity test (chi-squared) in the bivariate analysis to study inter-group differences. The dependent variable was dichotomic and distinguished between traffic offenders and drivers with no record. Median and standard deviations were calculated for quantitative variables, comparing groups making use of t-test. Size effect was calculated with Cohen's d .

To describe the traffic offenders' psychological profile, a binomial logistic regression was carried out, and a decision tree was derived using the CHAID method (Chi-squared Automatic Interaction Detector). CHAID is a hierarchical and divisive descending classification method, which aims to find group cases that may present similar and different characteristics with respect to other groups. In this investigation, it was intended to study

the segments that, using the Chi-Squared statistic, had a specific criterion which contrasted the possible differences between the segments formed through the interaction of the chosen independent variables (Escobar, 1998).

For the binomial logistic regression, all the predictor variables were introduced into the model as dummy variables (except for age, taken as a quantitative variable). All of them were nominal and dichotomic (the possibility of suffering a disorder or not). The conditions to apply this model were studied by analysing the normality of residues and the presence of the influential values that led to their overestimation by Cook's distance, to find that four subjects resulted in model overfit. Hence, in the end they were not included as participants, and the analyses continued with a final sample of 296 subjects. The forward Wald method was used for the selection of variables in the logistic regression model. The model was executed with 80 % of the 296 subjects (training sample with $n = 237$), of whom 36.16% did not commit offences ($n = 86$) and 63.83% were traffic offenders ($n = 151$). External validity was tested (test sample with $n = 59$) by the model's predictive capacity over the remaining 20.00%. The CHAID model was built with an 80% training sample, and its external validity was tested over the remaining 20%.

Statistical analyses were carried out using version 24.0 of the Statistical Package for Social Sciences (IBM SPSS). For this investigation, an .05 alpha level was used.

Results

Differences in presenting signs of PD, alcohol consumption and ADHD

Traffic offenders had higher scores than non-offenders in the general score on the Adult ADHD Self-Report Scale ($p = .027$) and in hyperactivity ($p < .001$), but not on the attention subscale. The effect size was small for ADHD, with total score ($d = 0.27$), and medium for hyperactivity ($d = 0.51$) (Table 2). On the other hand, the analysis revealed differences in frequencies between traffic offenders and non-offenders in presenting signs of PD, ADHD, and problematic alcohol use (Table 3).

A significant Chi-squared value appeared for the following: PDs, ordered according to level of significance: a) antisocial ($\chi^2 = 63.68$; $p < .001$), impulsive ($\chi^2 = 16.37$; $p < .001$) and borderline ($\chi^2 = 13.48$; $p < .001$); b) paranoid ($\chi^2 = 11.02$; $p = .001$) and anxious ($\chi^2 = 9.07$; $p = .003$). The results revealed that traffic offenders showed a higher frequency for the five PDs, where antisocial, impulsive, and borderline stood out. For ADHD, the statistics showed no statistically-significant differences between the groups, with 11% of traffic offenders presenting signs of ADHD. Statistically-significant differences appeared for problematic alcohol use ($\chi^2 = 46.72$; $p < .001$). Therefore, 55.80% of traffic offenders may have problematic alcohol consumption, far ahead of the figure for drivers with no traffic offences (13.90%) ($p < .001$).

Table 2.

Mean, standard deviation and statistical comparison for independent measures (specific factors of ADHD test).

ADHD	Traffic offenders (n=199)		Non-offender drivers (n=101)		<i>t</i>	<i>p</i> -value	<i>d</i>
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>			
General Score	7.98	3.72	6.99	3.56	2.22*	.027	0.27
Attention-Deficit	4.12	2.71	4.01	2.91	.30	.767	0.04
Hyperactivity	3.87	1.90	2.97	1.64	4.23***	.000	0.51

Note: *M*: Mean; *SD*: Standard Deviation; *t*: t-test; *d*: Cohen's *d*.

* = $p < .05$; ** = $p < .01$; *** = $p < .001$

Table 3.

Distribution of the frequencies and χ^2 of the variables of personality disorders, ADHD and problematic alcohol consumption.

Variable	Traffic offenders (n=199)		Non-offender drivers (n=101)		χ^2	<i>p</i> -value	<i>Phi</i>
	<i>n</i>	%	<i>n</i>	%			
Paranoid disorder	151	75.90	57	56.40	11.02**	.001	.20
Schizoid disorder	179	89.90	91	90.10	.00	1.000	.00
Antisocial disorder	145	72.90	24	23.80	63.68***	.000	.47
Impulsive disorder	58	29.10	8	7.90	16.37***	.000	.24
Borderline disorder	56	28.10	9	8.90	13.48***	.000	.22
Histrionic disorder	59	29.60	19	18.80	1.96	.060	.12
Anancastic disorder	147	73.90	66	65.30	1.97	.161	.09
Anxious disorder	107	53.80	35	34.70	9.07**	.003	.18
Dependent disorder	59	29.60	20	19.80	2.86	.091	.11
Without personality disorder	17	8.50	15	14.90	2.18	.140	.10
ADHD	22	11.10	5	5.00	2.35	.125	.10
Problematic alcohol consumption	111	55.80	14	13.90	46.72***	.000	.40

Note: χ^2 : Chi-squared Yates continuity correction. *Phi*: Phi Coefficient

* = $p < .05$; ** = $p < .01$; *** = $p < .001$.

Binomial regression logistics

As a result of applying the forward Wald method to select variables, the extracted model was formed by the predictor variables of gender, alcohol use-related problems, and borderline/antisocial PDs. Table 4 includes the model that resulted from Wald’s test to select variables. The last column shows the Beta exponent, which enables the interpretation of the odds ratio (OR). It can be concluded that alcohol use-related problems increased the probability of being convicted for traffic offences, with 9.5 traffic offenders in such circumstances for each driver with no such background. The presence of antisocial disorder increased the probability of being convicted for traffic offenders, with 7.6 offenders with signs of this disorder for each driver with no such background. The presence of borderline disorder increased the probability of being convicted for traffic offences, with 3.4 offenders with signs of this disorder for each driver with no such background. Lastly, there were 5.3 male traffic offenders for each female.

Table 4. Results of the logistic regression model for the traffic offender probability.

Variables	β	SE	Wald	gl	Sig.	Exp(β)
Gender (masculine)	1.672	.477	12.301	1	.000	5.323
Alcohol consumption disorder	2.261	.455	24.681	1	.000	9.592
Antisocial disorder	2.040	.380	28.778	1	.000	7.688
Borderline disorder	1.228	.600	4.182	1	.041	3.414
Constant	-2.580	.488	28.017	1	.000	.076

The model’s goodness of fit was tested using the Hosmer-Lemeshow test ($\chi^2 = 1.253, gl = 6, p = .974$). A good fit appeared with the observed values. The model’s goodness was also explored by studying the percentage of correctly-classified cases in the training sample, which came to 82.60%. Finally, the model’s predictive validity was found to be high, and correctly classified 84.70% of the remaining sample.

Decision tree

The CHAID method, employed with the training sample of 239 cases, reported how antisocial PD and alcohol consumption problems established significant differences in the probability of classifying someone as a traffic offender. The tree diagram (Figure 1) presents segmentation, which means that the most discriminating variable in the classification of a driver as a traffic offender is antisocial disorder. A higher probability of classifying a subject as a traffic offender (with 92.60%) was reflected among those subjects with signs of antisocial disorder and alcohol use problems.

The CHAID model was able to correctly classify 72.80% of the whole training sample (n = 239) and classified 78.70% of the reserve sample (n = 61) when testing its predictive validity.

Discussion

Different studies have analysed the relationship between risky conduct while driving and personality (Constantinou et al., 2011; Martí-Belda et al., 2019), antisocial conduct (Butcher et al., 2015; Gallardo & Andrés, 2009; Shechory et al., 2011) or alcohol abuse (Hilterman et al., 2012); which are predictor variables of its recidivism (Gugliotta, 2018; Herraiz et al., 2011; Monrás et al.,

2011). A higher prevalence has been found for ethylic dependence in people convicted of road safety crimes (Fáilde-Garrido et al., 2016; Valero, et al., 2017). In line with this, ADHD has been considered a neurodevelopmental disorder that is associated with antisocial (Simon, et al., 2009) and criminal (Pérez et al., 2015) behaviour, as has substance dependence (Andreu et al., 2015), because they can lead to these problems appearing in the life cycles of those suffering from them.




Figure 1. Traffic offender decision tree.

The present study aims to determine which psychological variables, from among those studied, are risk factors that can potentially impact road crimes. These results conclude that the most relevant variables are antisocial disorder and alcohol use problems, as well as hyperactivity. This finding concurs with previous related evidence (Keating et al., 2019) and confirms the research hypothesis. Additionally, a higher prevalence of different PDs (evaluated from a categorial perspective) has been reported in the group of traffic offenders, especially those related to lack of inhibitory control (borderline, antisocial, impulsive) (Albert et al., 2019, Broche-Pérez et al., 2016). This cognitive problem is also common in ADHD (particularly in the hyperactive subtype) (Ramos-Galarza et al., 2019), and can act as a shared element in all these clinical problems (Miranda et al., 2014), which might explain its relation to road crimes.

This study shows that, from ADHD symptoms, hyperactivity seems to be the most relevant for understanding this association, while attention remains stable between the groups and would not contribute to it. Hyperactivity has common traits among some personality disorders, such as impulsivity, which may suggest dimensional features that should be considered in the future.

Other dimensions, such as neuroticism and aggressiveness, might also be a point of convergence for various disorders and traffic offences (Slavinskienė et al., 2015).

The temporal-situational stability attributable to the personality structure (De Miguel & Pelechano, 2000) can, in turn, explain that the disorders in this area are associated with a high traffic offence relapse rate (Warren-Kigenyi & Coleman, 2014). Their therapeutic approach could hence be very important as a preventive measure. This is also in line with Knecht et al., (2015), who conclude that the risk of criminal behaviour among individuals with ADHD increases when there is a psychiatric comorbidity - specifically, a conduct disorder and a substance use disorder; therefore, these findings show the need to approach more individualised and comprehensive interventions, combining medical and psychosocial measures.

Data concerning the high prevalence of alcohol abuse are described herein, and concur with the data that other authors have previously reported (Hilterman et al., 2012). This abuse affects more than half of the sample, and comes over as one of the factors with the greatest classification capacity among groups. This finding is in line with the literature, which suggests the marked presence of alcoholism in people charged with road crimes (Alonso et al., 2019), and is most important because of its association with high mortality (Le Lièvre et al., 2019). In the current study, the concurrence of alcohol abuse and antisocial personality involves a model that suitably discriminates among groups. This backs existing pieces of evidence for the separate role of both (Cavaiola, 2003; Shechory et al., 2011), and also for alcohol consumption disorders along with reckless driving as a predictor of recidivism (Padilla et al., 2018). All this opens up new lines of research around the potential importance of their synergy. As previously considered for PDs and neurodevelopment, the depressor effect of alcohol on different executive functions (e.g., inhibitory control) can increase the risk of careless behaviour, especially when the personality structure that implies certain vulnerability to performing impulsive actions comes into play. Some pieces of evidence also exist for careful driving styles and a high perception of risk lowering the probability of committing offences while driving. These variables are compromised in ethylic intoxication (Padilla et al., 2018), and empower the tendency to inhibit responsibilities in antisocial disorders (APA, 2013).

A lack of significant results when comparing subjects in the presence of the clinical expression of ADHD suggests the need to explore which ADHD symptoms might be closely related to road safety crimes, by distinguishing a dimensional level (and not essentially a categorical one) in this matter. Future studies should evaluate the effect of attention deficit and hyperactivity separately, and how both are related to other mental disorders and impact the risk of committing traffic offences. More research is necessary to determine if psychiatric comorbidity in first-time offenders can predict the recidivism of road safety crimes. If this were the case, designing new assessment protocols that could identify these risk factors would be fundamental for assigning all the resources needed to impact them and to promote greater road safety. For example, reviewing and modifying the protocols for obtaining and renewing driving licences at driver assessment centres, which ensure the psychophysical conditions of people who drive as required by regulations.

Road safety is a first-order social problem because the accidents that occur while driving are one of the main causes of death in many countries around the world. Contributing to knowledge about the psychological variables that lie at its foundations will lead to efficient and evidence-based strategies. This article goes into detail about certain PDs as risk factors, and also about alcohol abuse, by providing evidence for them both sharing a higher prevalence in those subjects who commit road safety crimes. In view of the high comorbidity in these mental health problems, studying how they all influence, either together or separately, the complex equation of conduct while driving, and determining which variables are common to them all with a higher predictive power, is important.

Acknowledgements

The authors wish to thank the Deputy Directorate-General of Open Regime and Alternative Penal Measures, General Secretariat for Penitentiary Institutions (Ministry of the Interior), for authorising the participation of TASEVAL users for this study. They are also grateful for the support of its professionals.

Funding

This research received no financial aid from public sector agencies or commercial sector or not-for-profit organisations.

References

- Alavi, S.S., Mohammadi, M.R., Souri, H., Kalhori, S.M., Jannatifard, F. & Sepahbodi, G. (2017). Personality, Driving Behavior and Mental Disorders Factors as Predictors of Road Traffic Accidents Based on Logistic Regression. *Iranian Journal of Medical Sciences*, 42(1), 24-31. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5337762/>
- Albert, J., López-Martín, S., Arza, R., Palomares, N., Hoyos, S., Carretié, L., Díaz-Marsá, M., & Carrasco, J.L. (2019). Response Inhibition in Borderline Personality Disorder: Neural and Behavioral Correlates. *Biological Psychology*, 143, 32-40. <https://doi.org/10.1016/j.biopsycho.2019.02.003>
- Alonso, F., Esteban, C., Calatayud, C., Pastor, J. C., Alamar, B., & Medina, J. E. (2007). *La conducta social en el tráfico. Fundamentos para la intervención* [Social behavior in traffic. Rationale for intervention]. Attitudes. <https://hdl.handle.net/10550/50963>
- Alonso, F., Esteban, C., Sanmartín, J., & Useche, S. A. (2017). Reported prevalence of health conditions that affect drivers. *Cogent Medicine*, 4(1), Article 1303920. <https://doi.org/10.1080/2331205X.2017.1303920>
- Alonso, F., Esteban, C., Serge, A., & Tortosa, M. (2019). Importance of social and health-related problems: Do spaniards give them the significance they actually deserve? *International Journal of Environmental Research and Public Health* 16(21), Article 4090. <https://doi.org/10.3390/ijerph16214090>
- Alonso, F., Pastor, J. C., Montoro, L., & Esteban, C. (2015). Driving under the influence of alcohol: frequency, reasons, perceived risk and punishment. *Substance Abuse Treatment, Prevention, and Policy*, 10(1), 1-9. <https://doi.org/10.1186/s13011-015-0007-4>
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders (5th Ed.)*. <https://doi.org/10.1176/appi.books.9780890425596>

- Andreu, C., Letosa, J., López, M., & Mínguez, V. (2015). Implicaciones forenses en adultos con trastorno por déficit de atención con hiperactividad [Forensic implications in adults with attention deficit hyperactivity disorder]. *Revista Española de Medicina Legal*, 41(2), 65-71. <https://doi.org/10.1016/j.reml.2014.12.003>
- Ayinde, O. O., Adejumo, O. A., Olukolade, O., & Lasebikan, V. O. (2019). Should Traffic Offenders Undergo Compulsory “Mental Test”: A Study of Mental Health and Crash Involvement Among Commercial Motorcyclists in Ibadan, Nigeria? *Community Mental Health Journal*, 55(1), 180-187. <https://doi.org/10.1007/s10597-018-0302-4>
- Ballester, L., Alayo, I., Vilagut, G., Almenara, J., Cebrià, A. I., Echeburúa, E., Gabilondo, A., Gili, M., Lagares, C., Piqueras, J.A., Roca, M., Soto-Sanz, V., Blasco, M.J., Blasco, J., Castellví, P., Forero, C.G., Mortier, P., Alonso, J., & UNIVERSAL Study Group. (2021). Validation of an online version of the Alcohol Use Disorders Identification Test (AUDIT) for alcohol screening in Spanish university students. *International Journal of Environmental Research and Public Health*, 18(10), Article 5213. <https://doi.org/10.3390/ijerph18105213>
- Begg, D., Brookland, R., & Connor, J. (2017). Associations of repeated high alcohol use with unsafe driving behaviors, traffic offenses, and traffic crashes among young drivers: Findings from the New Zealand. *Traffic Injury Prevention*, 18(2), 111-117. <https://doi.org/10.1080/15389588.2016.1224345>
- Biçaksız, P., & Özkan, T. (2016). Impulsivity and driver behaviors, offences and accident involvement: A systematic review. *Transportation Research Part F: Traffic Psychology and Behaviour*, 38, 194-223. <https://doi.org/10.1016/j.trf.2015.06.001>
- Broche-Pérez, Y., & Herrera-Jiménez, L.F. (2016). Funciones ejecutivas “frías” y “calientes” en adolescentes con trastorno disocial [Executive functions in adolescents with conduct disorder]. *Revista Neuropsicología, Neuropsiquiatría y Neurociencias*, 16(3), 157-175.
- Butcher, J. N., Hass, G. A., Greene, R. L., & Nelson, L. D. (2015). *Using the MMPI-2 in forensic assessment*. American Psychological Association.
- Cabasés, J., & Quintero, F. (2005). Tratamiento multimodal del trastorno por déficit de atención con hiperactividad (TDAH) [Multimodal treatment of attention deficit hyperactivity disorder (ADHD)]. *Revista Española de Pediatría*, 61(6), 503-514. <https://www.seinap.es/wp-content/uploads/Revista-de-Pediatría/2005/REP%2061-6.pdf>
- Calinescu, T., & Adnanaite, D. (2018). *Progress in reducing drink driving in Europe*. European Transport Safety Council. https://etsc.eu/wp-content/uploads/report_reducing_drink_driving_final.pdf
- Cavaiola, A.A., Strohmetz, D.B., Wolf, J.M., & Lavender, N.J. (2003). Comparison of DWI offenders with non-DWI individuals on the MMPI-2 and the Michigan Alcoholism Screening Test. *Addictive Behaviors*, 28(5), 971-977. [https://doi.org/10.1016/S0306-4603\(01\)00291-X](https://doi.org/10.1016/S0306-4603(01)00291-X)
- Chang, Z., Quinn, P., Hur, K., Gibbons, R., Sjölander, A., Larsson, H., & D’Onofrio, B. (2017). Association between medication use for attention-deficit/ hyperactivity disorder and risk of motor vehicle crashes. *JAMA Psychiatry*, 74(6), 597-603. <https://doi.org/10.1001/jamapsychiatry.2017.0659>
- Chen, T.Y. & Jou, R.C. (2018). Estimating factors of individual and regional characteristics affecting the drink driving recidivism. *Accident, Analysis & Prevention*, 119, 16-22. <https://doi.org/10.1016/j.aap.2018.06.011>
- Constantinou, E., Panayiotou, G., Konstantinou, N., Loutsiou-Ladd, A., & Kapardis, A. (2011). Risky and aggressive driving in young adults: Personality matters. *Accident Analysis & Prevention*, 43(4), 1323-1331. <https://doi.org/10.1016/j.aap.2011.02.002>
- Curry, A. E., Metzger, K. B., Pfeiffer, M. R., Elliott, M. R., Winston, F. K., & Power, T. J. (2017). Motor vehicle crash risk among adolescents and young adults with attention-deficit/hyperactivity disorder. *JAMA Pediatrics*, 171(8), 756-763. <https://doi.org/10.1001/jamapediatrics.2017.0910>
- Daigre, C., Ramos-Quiroga, J. A., Valero, S., Bosch, R., Roncero, C., Gonzalvo, B., Nogueira, M., & Casas, M. (2009). Cuestionario autoinformado de cribado de TDAH ASRS-v1. 1 en adultos en tratamiento por trastornos por uso de sustancias [Adult ADHD Self-Report Scale (ASRS-v1.1) symptom checklist in patients with substance use disorders]. *Actas Españolas de Psiquiatría*, 37(6), 299-305. <https://www.researchgate.net/publication/277272661>
- Dekkers, T. J., Popma, A., van Rentergem, J. A. A., Bexkens, A., & Huizenga, H. M. (2016). Risky decision making in attention-deficit/hyperactivity disorder: A meta-regression analysis. *Clinical Psychology Review*, 45, 1-16. <https://doi.org/10.1016/j.cpr.2016.03.001>
- De las Cuevas, C., & Sanz, E.J. (2009). Enfermedad mental y seguridad vial [Mental illness and road safety]. *Actas Españolas de Psiquiatría*, 37(2), 75-81. <https://medes.com/publication/51940>
- De Miguel, A., & Pelechano, V. (2000). Estructura de personalidad y trastornos de personalidad: Correlatos de personalidad de los trastornos de personalidad definidos por clasificaciones de consenso [Personality structure and personality disorders: Personality correlates of personality disorders defined by consensus classifications]. *Psicologemas*, 14(27-28), 1-210. <https://dialnet.unirioja.es/servlet/revista?codigo=12352>
- Del Río, M. C., Gómez, J., Sancho, M., & Álvarez, F. J. (2002). Alcohol, illicit drugs and medicinal drugs in fatally injured drivers in Spain between 1991 and 2000. *Forensic Science International*, 127(1-2), 63-70. [https://doi.org/10.1016/S0379-0738\(02\)00116-0](https://doi.org/10.1016/S0379-0738(02)00116-0)
- Del Río, M. C., González-Luque, J. C., & Álvarez, F. J. (2001). Alcohol-related problems and fitness to drive. *Alcohol & Alcoholism*, 36(3), 256-261. <https://doi.org/10.1093/alcalc/36.3.256>
- Directive 2006/126/EC of the European Parliament and of the Council of 20 December 2006 on Driving Licences (Recast) (Text with EEA Relevance). *The Official Journal of the European Union*, L403, 18-60, of 30 December 2006. <https://data.europa.eu/eli/dir/2006/126/oj>
- Directorate General of Traffic (2020). *Las principales cifras de la siniestralidad vial. España, 2019* [The main figures of road accidents. Spain, 2019]. Dirección General de Tráfico. <https://www.dgt.es/menusecundario/dgt-en-cifras/dgt-en-cifras-resultados/dgt-en-cifras-detalle/?id=00151>
- Ensoo, D., Kiive, E., Laas, K. & Harro, J. (2015). The effect of ADHD symptoms in traffic accidents. *Journal of Local and Global Health Science*, 2, 8. <https://doi.org/10.5339/jlghs.2015.itma.8>
- Escobar, M. (1998). Las aplicaciones del análisis de segmentación: el procedimiento Chaid [Applications of segmentation analysis: The Chaid procedure]. *EMPIRIA. Revista de Metodología de Ciencias Sociales*, 1, 13-49. <https://doi.org/10.5944/empiria.1.1998.706>
- European Commission (June 11, 2020). *2019 road safety statistics: what are the numbers behind?* https://ec.europa.eu/commission/presscorner/detail/en/qanda_20_1004
- Failde-Garrido, J. M., García-Rodríguez, M. A., Rodríguez-Castro, Y., González-Fernández, A., Fernández, M. L., & Fernández, M. V. C. (2016). Psychosocial determinants of road traffic offences in a sample of Spanish male prison inmates. *Transportation Research Part F: Traffic Psychology and Behaviour*, 37, 97-106. <https://doi.org/10.1016/j.trf.2015.12.004>

- Freeman, J., Maxwell, J.C., & Davey, J. (2011). Unraveling the complexity of driving while intoxicated; A study into the prevalence of psychiatric and substance abuse comorbidity. *Accident Analysis and Prevention*, 43(1), 34-39. <https://doi.org/10.1016/j.aap.2010.06.004>
- Gallardo, D., & Andrés, A. (2009). Perfil de un grupo de penados por delitos contra la seguridad del tráfico [Profile of a group of convicts for crimes against traffic safety]. *Invesbren Criminología*, 48, 2-6. <https://dialnet.unirioja.es/ejemplar/244936>
- García, T., Rodríguez, C., Rodríguez, J., Fernández-Suárez, A., Richarte, V., & Ramos-Quiroga, J. A. (2019). Psychosocial profiles of adults with ADHD: A comparative study of prison and outpatient psychiatric samples. *The European Journal of Psychology Applied to Legal Context*, 11(1), 41-49. <https://doi.org/10.5093/ejpalc2018a14>
- General State Prosecutor's Office (2020). *Memoria 2020* [Memory 2020]. Fiscalía General del Estado. Ministerio de Justicia. https://www.fiscal.es/memorias/memoria2020/FISCALIA_SITE/index.html
- Gómez-Talegón, M. T., Fierro, I., Vicondoa, Á., Ozcoide, M., & Álvarez, F. J. (2008). Aptitud para conducir de los pacientes con trastornos mentales. Su evaluación en los centros de reconocimiento de conductores [Fitness to drive among patients with mental disorders. Their assessment in medical centers for drivers]. *Psiquiatría Biológica*, 15(3), 63-72. [https://doi.org/10.1016/S1134-5934\(08\)71125-4](https://doi.org/10.1016/S1134-5934(08)71125-4)
- González-Iglesias, B., & Gómez-Fraguela, J. A. (2010). Conductores infractores, ¿un perfil de conducta desviada? Análisis de las diferencias y similitudes con una muestra de conductores de la población general [Violating drivers, a deviant behavior profile? Analysis of differences and similarities with a sample of drivers from the general population]. *Revista Española de Investigación Criminológica*, 8, 1-19. <https://doi.org/10.46381/reic.v8i0.56>
- Gugliotta, A.A. (2018). *Análisis de la percepción de peligros y la estimación del riesgo para definir un perfil del conductor seguro* (Publicación No. 9788491637868) [Hazard perception analysis and risk estimation to define a safe driver profile] [Doctoral dissertation, University of Granada]. UG Campus Repository. <https://hdl.handle.net/10481/49718>
- Harpin, V.A. (2005). The effect of ADHD on the life of an individual, their family, and community from preschool to adult life. *Archives Disease in Childhood*, 90(1), 2-7. <https://dx.doi.org/10.1136/adc.2004.059006>
- Herraiz, F. (2009). Descripción del perfil psicológico de los internos por delitos contra la seguridad del tráfico [Description of the psychological profile of inmates for crimes against traffic safety]. *Invesbren Criminología*, 48, 9-11. <https://dialnet.unirioja.es/ejemplar/244936>
- Herraiz, F., Chamorro, A., & Villamarín, F. (2011). Predictores psicosociales de delitos contra la seguridad vial: un estudio con internos de un centro penitenciario. [Psychosocial predictors of traffic crimes: A study with inmates in a penitentiary centre]. *Clínica y Salud*, 22(2), 87-100. <https://doi.org/10.5093/cl2011v22n2a1>
- Hilterman, E., Mancho, R., Trasovares, M.V., Baldris, G., & Muñoz, Y. (2012). Programas de tratamiento y características de los internos penitenciarios ingresados por delitos de tráfico en Catalunya [Treatment programs and characteristics of prison inmates admitted for traffic offenses in Catalunya]. *Invesbren Criminología*, 57, 6-11. <https://dialnet.unirioja.es/servlet/articulo?codigo=6273037>
- Jiménez- Arriero, J., Rodríguez-Jiménez, R., Vidal de la Fuente, J., & Ponce Alfaro, G. (2005). El TDAH: Evolución a la edad adulta [ADHD: Evolution to adult age]. *Revista Española de Pediatría*, 61(6), 495-500. <https://www.seinap.es/wp-content/uploads/Revista-de-Pediatría/2005/REP%2061-6.pdf>
- Jonah, B.A., Thiessen, R., & Au-Yeung, E. (2001). Sensation seeking, risky driving and behavioral adaptation. *Accident Analysis & Prevention* 33(5), 679-684. [https://doi.org/10.1016/S0001-4575\(00\)00085-3](https://doi.org/10.1016/S0001-4575(00)00085-3)
- Jornet-Gibert, M., Gallardo-Pujol, D., Suso, C., & Andrés-Pueyo, A. (2013). Attitudes do matter: The role of attitudes and personality in DUI offenders. *Accident Analysis & Prevention*, 50, 445-450. <https://doi.org/10.1016/j.aap.2012.05.023>
- Kaye, S., Gilseman, J., Young, J. T., Carruthers, S., Allsop, S., Degenhardt, L., ... & van den Brink, W. (2014). Risk behaviours among substance use disorder treatment seekers with and without adult ADHD symptoms. *Drug and Alcohol Dependence*, 144, 70-77. <https://doi.org/10.1016/j.drugalcdep.2014.08.008>
- Keating, L. M., Nelson, S. E., Wiley, R. C., & Shaffer, H. J. (2019). Psychiatric comorbidity among first-time and repeat DUI offenders. *Addictive Behaviors*, 96, 1-10. <https://doi.org/10.1016/j.addbeh.2019.03.018>
- Kessler, R. C., Adler, L., Ames, M., Delmer, O., Faraone, S., Hiripi, E., Howes, M. J., Jin, R., Secnik, K., Spencer, T., Ustun, T. B., & Walters, E. E. (2005). The World Health Organization adult ADHD self-report scale (ASRS): A short screening scale for sse in the general population. *Psychological Medicine*, 35(2), 245-256. <https://doi.org/10.1017/S0033291704002892>
- Kessler, R. C., Adler, L. A., Gruber, M. J., Sarawate, C. A., Spencer, T., & Van Brunt, D. L., (2007). Validity of the World Health Organization adult ADHD self-report scale (ASRS) screener in a representative sample of health plan members. *International Journal of Methods in Psychiatric Research*, 16(2), 52-65. <https://doi.org/10.1002/mpr.208>
- Kieling, R. R., Szobot, C. M., Matte, B., Coelho, R. S., Kieling, C., Pechansky, F., & Rohde, L. A. (2011). Mental disorders and delivery motorcycle drivers (motoboy): a dangerous association. *European Psychiatry*, 26(1), 23-27. <https://doi.org/10.1016/j.eurpsy.2010.03.004>
- Klein, R. G., Mannuzza, S., Olazagasti, M. A. R., Roizen, E., Hutchison, J. A., Lashua, E. C., & Castellanos, F. X. (2012). Clinical and functional outcome of childhood attention-deficit/hyperactivity disorder 33 years later. *Archives of General Psychiatry*, 69(12), 1295-1303. <https://doi.org/10.1001/archgenpsychiatry.2012.271>
- Knecht, C., de Alvaro, R., Martínez-Raga, J., & Balanza-Martínez, V. (2015). Attention-deficit hyperactivity disorder (ADHD), substance use disorders, and criminality: a difficult problem with complex solutions. *International journal of adolescent medicine and health*, 27(2), 163-175. <https://doi.org/10.1515/ijamh-2015-5007>
- Landis, J. R., & Koch, G. G. (1977). The measurement of observer agreement for categorical data. *Biometrics*, 33(1), 159-174. <https://doi.org/10.2307/2529310>
- Lapham, S. C, Smith, E., C' de Baca, J., Chang, I., Skipper, B.J., Baum, G., & Hunt, W.C. (2001). Prevalence of psychiatric disorders among persons convicted of driving while impaired. *Archives of General Psychiatry*, 58(10), 943-949. <https://doi.org/10.1001/archpsyc.58.10.943>
- Le Lièvre, P., Admindaite, D., Jost, G., & Podda, F. (2019). *Progress in reducing drink-driving and other alcohol-related road deaths in Europe*. European Transport Safety Council. <https://etsc.eu/progress-in-reducing-drink-driving-in-europe-2019/>
- López, J.A., Serrano, I., & Delgado, J. (2005). Attention deficit hyperactivity disorder: a predictive model of comorbidity with behaviour disorder. *Psychology in Spain*, 9(1), 63-74. <https://www.psychologyinspain.com/content/full/2005/9008.pdf>
- Loranger A.W., Janca A., & Sartorius N. (Eds.) (1997). *Assessment and diagnosis of personality disorders: the ICD-10 international personality disorder examination (IPDE)*. Cambridge University Press. <https://apps.who.int/iris/handle/10665/41912>

- Luderer, M., Quiroga, J. A. R., Faraone, S. V., Zhang-James, Y., & Reif, A. (2021). Alcohol use disorders and ADHD. *Neuroscience & Biobehavioral Reviews*, 128, 648-660. <https://doi.org/10.1016/j.neubiorev.2021.07.010>
- Marti-Belda, A., Pastor, J. C., Montoro, L., & Bosó, P. (2019). Persistent traffic offenders: Alcohol consumption and personality as predictors of driving disqualification. *The European Journal of Psychology Applied to Legal Context*, 11(2), 81-92. <https://doi.org/10.5093/ejpalc2019a3>
- McCutcheon, V. V., Heath, A. C., Edenberg, H. J., Grucza, R. A., Hesselbrock, V. M., Kramer, J. R., Bierut, L. J., & Bucholz, K. K. (2009). Alcohol criteria endorsement and psychiatric and drug use disorders among DUI offenders: Greater severity among women and multiple offenders. *Addictive Behaviors*, 34(5), 432-439. <https://doi.org/10.1016/j.addbeh.2008.12.003>
- McDonald, A. S., & Davey, G. C. (1996). Psychiatric disorders and accidental injury. *Clinical Psychology Review*, 16(2), 105-127. [https://doi.org/10.1016/0272-7358\(96\)00007-4](https://doi.org/10.1016/0272-7358(96)00007-4)
- Miranda, A., Berenguer, C., Colomer, C., & Roselló, R. (2014). Influence of the symptoms of Attention Deficit Hyperactivity Disorder (ADHD) and comorbid disorders on functioning in adulthood. *Psicothema*, 26(4), 471-476. <https://doi.org/10.7334/psicothema2014.121>
- Monrás, M., Aparicio, Á., López, J. A., & Pons, I. (2011). Causas de los ingresos en prisión por delitos contra la seguridad vial: ¿Enfermedad o irresponsabilidad? [Causes of imprisonment for crimes against road safety: Illness or irresponsibility?] *Medicina Clínica*, 137(6), 279-280. <https://doi.org/10.1016/j.medcli.2010.07.012>
- Moskowitz, H., & Fiorentino, D. A. (2000). *Review of the Literature on the Effects of Low Doses of Alcohol on Driving-Related Skills*. National Highway Traffic Safety Administration. <https://one.nhtsa.gov/people/injury/research/pub/hs809028/Title.htm>
- National Institute of Toxicology and Forensic Sciences (2020). *Hallazgos toxicológicos en víctimas mortales de accidentes de tráfico. Memoria 2019* [Toxicological findings in traffic accident fatalities. Memory, 2019]. Ministerio de Justicia. https://www.mjusticia.gob.es/es/ElMinisterio/OrganismosMinisterio/Documents/1292430960214-Memoria_Trafico_INTCF_2019.PDF
- Nelson, S. E., Belkin, K., LaPlante, D. A., Bosworth, L., & Shaffer, H. J. (2015). A prospective study of psychiatric comorbidity and recidivism among repeat DUI offenders. *Archives of Scientific Psychology*, 3(1), 8-17. <https://dx.doi.org/10.1037/arc0000009>
- Nelson, S. E., Shoov, E., LaBrie, R. A., & Shaffer, H. J. (2019). Externalizing and self-medicating: Heterogeneity among repeat DUI offenders. *Drug and Alcohol Dependence*, 194, 88-96. <https://doi.org/10.1016/j.drugalcdep.2018.09.017>
- Nordfjaern, T. & Şimşekoğlu, Ö. (2014). Empathy, conformity, and cultural factors related to aberrant driving behaviour in a sample of urban Turkish drivers. *Safety Science*, 68, 55-64. <https://doi.org/10.1016/j.ssci.2014.02.020>
- Padilla, J. L., Doncel, P., Gugliotta, A., & Castro, C. (2018). Which drivers are at risk? Factors that determine the profile of the reoffender driver. *Accident Analysis & Prevention*, 119, 237-247. <https://doi.org/10.1016/j.aap.2018.07.021>
- Pérez, C. R., Pérez, J. C. N., Díaz, F. J. R., Granda, A. P., Molleda, C. B., & Fernández, T. G. (2015). Trastorno por Déficit de Atención e Hiperactividad (ADHD): Prevalencia y características sociodemográficas en población reclusa [Attention Deficit Hyperactivity Disorder (ADHD): Prevalence and sociodemographic features in imprisoned population]. *Psicología Reflexão e Crítica*, 28(4), 698-707. <https://doi.org/10.1590/1678-7153.201528407>
- Pérola de Torres, L. P., Fernández-García, J. A., Arias-Vega, R., Muriel-Palomino, M., Márquez-Rebollo, E., & Ruiz-Moral, R. (2005). Validación del cuestionario AUDIT para la identificación del consumo de riesgo y de los trastornos por el uso de alcohol en mujeres [Validation of the AUDIT test for identifying risk consumption and alcohol use disorders in women]. *Atención Primaria*, 36(9), 499-506. [https://doi.org/10.1016/S0212-6567\(05\)70552-7](https://doi.org/10.1016/S0212-6567(05)70552-7)
- Räsänen, T., Hakko, H., Riipinen, P., Rätty, E. & Kantojärvi, L. (2019). Personality disorders of drivers killed in fatal motor vehicle accidents in Finland during 1990–2011. *Acta Psychiatrica Scandinavica*, 140(1), 39-49. <https://doi.org/10.1111/acps.13039>
- Ramos-Galarza, C., Acosta-Rodas, P., Pérez-Salas, C., & Ramos, V. (2019). Inhibitory control and symptomatology of Attention Deficit Hyperactivity Disorder. *Revista Ecuatoriana de Neurología*, 28(3), 41-46. <https://scielo.senescyt.gob.ec/pdf/rneuro/v28n3/2631-2581-rneuro-28-03-00041.pdf>
- Ramos-Quiroga, J. A., Bosch-Munsó, R., Castells-Cervelló, X., Nogueira-Morais, M., García-Giménez, E., & Casas-Brugué, M. (2006). Trastorno por déficit de atención con hiperactividad en adultos: Caracterización clínica y terapéutica [Attention deficit hyperactivity disorder in adults: A clinical and therapeutic characterization]. *Revista de Neurología*, 42(10), 600-606. <https://doi.org/10.33588/rn.4210.2005495>
- Ramos-Quiroga, J. A., Montoya, A., Kutzelnigg, A., Deberdt, W., & Sobanski, E. (2013). Attention deficit hyperactivity disorder in the European adult population: prevalence, disease awareness, and treatment guidelines. *Current Medical Research and Opinion*, 29(9), 1093-1104. <https://doi.org/10.1185/03007995.2013.812961>
- Retz, W., Ginsberg, Y., Turner, D., Barra, S., Retz-Junginger, P., Larsson, H., & Asherson, P. (2021). Attention-Deficit/Hyperactivity Disorder (ADHD), antisociality and delinquent behavior over the lifespan. *Neuroscience & Biobehavioral Reviews*, 120, 236-248. <https://doi.org/10.1016/j.neubiorev.2020.11.025>
- Rubio, G., Bermejo, J., Caballero, M. C., & Santo-Domingo, J. (1998). Validación de la prueba para la identificación de trastornos por uso de alcohol (AUDIT) en Atención Primaria [Validation of the alcohol use disorders identification test (AUDIT) in primary care] *Revista Clínica Española*, 198(1), 11-14. <https://pubmed.ncbi.nlm.nih.gov/9534342/>
- Schell, T. L., Chan, K. S., & Morral, A. R. (2006). Predicting DUI recidivism: Personality, attitudinal, and behavioral risk factors. *Drug and Alcohol Dependence*, 82(1), 33-40. <https://doi.org/10.1016/j.drugalcdep.2005.08.006>
- Shechory, M., Weiss, J. M., & Weinstein, R. (2011). Differentiating offenders by index offense and personality inventories: The characteristics of adult probationers in Israel. *International Journal of Offender Therapy and Comparative Criminology* 57(3) 312–331. <https://doi.org/10.1177/0306624X11428316>
- Simon, V., Czobor, P., Bálin, S., Meszaros, A., & Bitter, I. (2009). Prevalence and correlates of adult attention-deficit hyperactivity disorder: Meta-analysis. *The British Journal of Psychiatry*, 194(3), 204-211. <https://doi.org/10.1192/bjp.bp.107.048827>
- Slavinskienė, J., Žardeckaitė-Matulaitienė, K., Markšaitytė, R., Šeibokaitė, L., & Endriulaitienė, A. (2015). Personality profiles of Lithuanian traffic offenders. *Transport means - 2015: Proceedings of the 19th international scientific conference, October 22-23, 2015, Kaunas University of Technology, Lithuania*, 19, 207-210. <https://hdl.handle.net/20.500.12259/30801>
- Soutullo, C., & Diez, A. (2007). *Manual de Diagnóstico y Tratamiento del TDAH* [ADHD Diagnosis and Treatment Manual]. Ed. Medica Panamericana.

- Spanish Organic Law 15/2007 (November 30, 2007). Modificación de la Ley Orgánica 10/1995, de 23 de noviembre, del Código Penal en materia de seguridad vial [Modification of Organic Law 10/1995, of November 23, of the Criminal Code on road safety] *Boletín Oficial del Estado*, 288, December 1, 2007. <https://www.boe.es/eli/es/lo/2007/11/30/15>
- Spanish Royal Decree, RD 818/2009 (May 8, 2009). Reglamento General de Conductores [General Drivers Regulation]. *Boletín Oficial del Estado*, 138, June 8, 2009. <https://www.boe.es/eli/es/rd/2009/05/08/818>
- Tortosa, F., & Montoro, L. (2002). La Psicología aplicada a la selección de conductores. Cien años salvando vidas [Psychology applied to the selection of drivers. One hundred years saving lives]. *Psicothema*, 14(4), 714-725. <https://www.psicothema.com/pii?pii=789>
- Trujillo, N., Pineda, D.A., & Puerta, I.C. (2007). Alteraciones cognitivas en adolescentes infractores con trastorno disocial de diversos niveles de gravedad [Cognitive alterations in offending adolescents with conduct disorder of various levels of severity]. *Psicología Conductual*, 15(2), 297-319. https://www.researchgate.net/publication/228369610_Alteraciones_cognitivas_en_adolescentes_infractores_con_trastorno_disocial_de_diversos_niveles_de_gravedad
- Vaa, T. (2013). ADHD and relative risk of accidents in road traffic: A meta-analysis. *Accident: Analysis and Prevention*, 62, 415-425. <https://doi.org/10.1016/j.aap.2013.10.003>
- Valdizán J. R. y Izaguerri-Gracia A. C. (2009). Trastorno por déficit de atención/hiperactividad en adultos [Attention deficit/hyperactivity disorder in adults]. *Revista de Neurología*, 48(Supl 2), S95-S99. <https://doi.org/10.33588/rn.48S02.2009017>
- Valero, S., Bosch, R., Corominas, M., Giannoni, A., Barrau, V., Ramos Quieroga, J. Al, & Casas, M. (2017). Psychopathology and traffic violations in subjects who have lost their driving license. *Comprehensive Psychiatry*, 76, 45-55. <https://doi.org/10.1016/j.comppsy.2017.03.012>
- Warren-Kingenyi, N., & Coleman, H. (2014). *DWI Recidivism in the United States: An examination of state-level driver data and the effect of look-back periods on recidivism prevalence*. National Highway Traffic Safety Administration. <https://trid.trb.org/view/1306862>
- Wijnen, W., & Stipdonk, H. (2016). Social costs of road crashes: An international analysis. *Accident Analysis & Prevention* 94, 97-106. <https://doi.org/10.1016/j.aap.2016.05.005>
- World Health Organization (1992). *International Statistical Classification of Diseases and Related Health Problems (ICD-10). Mental and behavioural disorders*.
- World Health Organization. (2001). *AUDIT: The Alcohol Use Disorders Identification Test: guidelines for use in primary health care*. 2nd ed. World Health Organization. <https://apps.who.int/iris/handle/10665/67205>
- World Health Organization (2003). *Self-Report Scale Version V1.1 (ASRS-v1.1) Symptoms Checklist*.
- World Health Organization (2018, June 17). *Global status report on road safety 2018*. https://www.who.int/violence_injury_prevention/road_safety_status/2018/en/